



11 Publication number: **0 371 802 B1**

12 **EUROPEAN PATENT SPECIFICATION**

43 Date of publication of patent specification :
11.11.92 Bulletin 92/46

51 Int. Cl.⁵: **A61K 7/48, A61K 7/06,**
A61K 7/50, A61K 7/08

21 Application number: 89312485.9

22 Date of filing: 30.11.89

54 **Topical composition.**

30 Priority: 01.12.88 GB 8828013

43 Date of publication of application :
06.06.90 Bulletin 90/23

45 Publication of the grant of the patent :
11.11.92 Bulletin 92/46

84 Designated Contracting States :
AT BE CH DE ES FR GB GR IT LI NL SE

56 References cited :
EP-A- 0 018 717
EP-A- 0 077 920
EP-A- 0 224 796
WO-A-89/01023

73 Proprietor: UNILEVER PLC
Unilever House Blackfriars P.O. Box 68
London EC4P 4BQ (GB)

84 GB
Proprietor: UNILEVER N.V.
Burgemeester s'Jacobplein 1 P.O. Box 760
NL-3000 DK Rotterdam (NL)

84 BE CH DE ES FR GR IT LI NL SE AT

72 Inventor: Birtwistle, David Howard
18 Exmoor Close Irby
Wirral Merseyside, L61 9QN (GB)
Inventor: Carter, Peter
Briarfield The Rake
Burton Wirral Merseyside (GB)
Inventor: Rosser, David Arthur
The Sycamores The Mount
Heswall Wirral Merseyside, L60 4RD (GB)

74 Representative: Elliott, Peter William et al
Unilever plc, Patent Division Colworth House
Sharnbrook
Bedford MK44 1LQ (GB)

EP 0 371 802 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

FIELD OF THE INVENTION

- 5 The invention relates to compositions suitable for topical application to the skin, (including the mucosae), and to the hair. In particular, the invention is concerned with highly improved detergent compositions suitable for cleansing the whole body surface, including the mouth.

BACKGROUND TO THE INVENTION & PRIOR ART

- 10 The damaging effect of conventional detergents used to wash the body surface, particularly where young, tender or damaged skin is involved, has been the subject of intense study for many years in a search for milder-to-the-skin products, which not only cleanse the skin efficiently, but also leave the skin with a pleasant smooth silky feel after the skin surface has been dried off.

- 15 The use of certain mono- and di-alkyl phosphate salts for this purpose has been advocated in view of their mild characteristics, but some of this group of salts are used as antifoam agents because of their lather suppressant properties, and would therefore require careful formulation if lather control is not required.

- To this end, US patent No. 4,139,485 (Kao Soap Co. Ltd.) describes a detergent composition having low irritation properties on human skin, wherein the surfactant component is dialkyl or dialkenyl phosphate salt (DAP) and/or monoalkyl or monoalkenyl phosphate salt (MAP), each alkyl or alkenyl group having from 10 to 20 16 carbon atoms), the weight ratio of 'DAP' to 'MAP' being from 20:80 to 0:100. This system is stated to possess good detergency.

- Also, US patent No. 4,526,710 (Kao Corporation), report a study to improve properties of detergent composition which make use of anionic phosphate surface active agents, which Kao maintain are highly innocuous and particularly mild to the skin. As a result, Kao have found that when phosphate ester salts having a specific ion pair, notably mono- or di-alkyl (C8-18) phosphates, are used in combination with alkanol amine salts of high- 25 fatty acids and alkyl amine oxides, the detergency and foaming characteristics are remarkably improved.

- Also, US patent No. 4,758,376 (Kao) discloses an alternative composition, comprising an alkanolamine salt of a mono or dialkyl (C8-18) phosphate or mixtures thereof, to that described in US patent No. 4,526,710 in which the problem of poor foaming due to the dialkyl phosphate salt is dealt with by incorporating with the phosphate a compound chosen from an amidoamine amphoteric surfactant or hydroxysulphobetaine, or an ali- 30 phatic lactylate or glycolate.


- It is clear from a study of the three Kao references, that all the evidence on which their disclosures are based is restricted to alkyl phosphates or dialkyl phosphates in which the alkyl group has at least 10 carbon atoms. There is absolutely no reference at all to any evaluation of an alkyl phosphate ester or dialkyl phosphate ester salt where the alkyl group or groups contains only 9 carbon atoms, or less.

- Applicants in their search for a mild surfactant for use in cleansing human skin or hair, with the added attribute that a full, soft lather is produced without necessarily incorporating a secondary surfactant such as those proposed by Kao, have unexpectedly discovered that certain dioctyl phosphate salts meet their requirements in this respect. Furthermore, even higher lather volumes can be produced when the amount to be incorporated in such formulations of the corresponding mono-octyl phosphate salt, which usually contaminates commercially available dioctyl phosphate salt, is reduced to a lower than usual level. The compositions so obtained are accordingly capable of producing a superior lather volume and an outstanding lather creaminess. Also, the composition is so mild to the skin that it can safely be used for cleansing the mucosae, such as the mouth and the vagina, and other more delicate skin areas. It can also be used in shampoos for frequent, e.g. daily, hair washing, without risk of scalp irritation or damage attributable to harsher products. In addition to these excellent attributes, the ease of rinsing from hair or skin and superior silky-smooth after-use skin feel properties of the compositions, including freedom from skin roughness and erythema, have great consumer appeal.


DEFINITION OF THE INVENTION

- 50 Accordingly, the invention provides a composition suitable for topical application to the skin or hair, which comprises:

- (a) from 1 to 99.9% by weight of a dialkyl or dialkenyl phosphate salt (or mixtures thereof) having the structure (1):



(b) from 0.01 to 20% by weight of a branched or unbranched, monoalkyl or monoalkenyl phosphate salt (or mixtures thereof) having the structure (2):



where R^{01} , and R^{02} and R^{03} are each chosen from branched or unbranched, alkyl or alkenyl groups having an average of from 7 to 9 carbon atoms;

X, Y and Z are each chosen from H, alkali metal, ammonium and substituted ammonium counterions; and


a, b and c are each chosen from 0 or a value of from 1 to 4;

the weight ratio of the monoalkyl or monoalkenyl phosphate salt to the dialkyl or dialkenyl phosphate salt being from 1:1000 to 1:5.

DISCLOSURE OF THE INVENTION

The Dialkyl or Dialkenyl Phosphate Salt

The composition according to the invention comprises as a surfactant, a dialkyl or dialkenyl phosphate salt (or mixtures thereof) having the structure:



where R^{01} and R^{02} are each chosen from branched or unbranched alkyl or alkenyl groups having an average of from 7 to 9 carbon atoms;

X is chosen from H, alkali metal ammonium, and substituted ammonium counterions; and

a & b are each chosen from 0 or a value of from 1 to 4.

Examples of the dialkyl- and dialkenyl phosphate moiety include:

di-n-heptyl phosphate

di-n-octyl phosphate

di-n-nonyl phosphate

di-(diethyleneglycol-mono-n-nonyl ether) phosphate

di-(ethyleneglycol-mono-n-octyl ether) phosphate

di-n-heptenyl phosphate

di-n-octenyl phosphate

di-n-nonenyl phosphate

di-(triethyleneglycol-mono-n-octenyl ether) phosphate

di-(ethyleneglycol-mono-n-heptyl ether) phosphate

di-(7-methyloctyl) phosphate

di-(5-methylhexyl) phosphate
 di-(6,6-dimethylheptyl) phosphate
 n-hexyl-n-nonyl phosphate, and
 n-hexyl-n-octyl phosphate.


5 The preferred dialkyl phosphate moiety is di-n-octyl phosphate, and the preferred counterion is chosen from sodium, potassium and triethanolammonium.

The amount of dialkyl or dialkenyl phosphate salt which is present in the composition according to the invention is from 1 to 99.9%, preferably from 2 to 50% by weight of the composition.

10 Compositions containing less than 1% by weight of the dialkyl or dialkenyl phosphate salt show poor lather characteristics and lack the desired feel properties associated with compositions containing higher levels of this ingredient.

The Monoalkyl or Monoalkenyl Phosphate Salt

15 The composition according to the invention also comprises, as a lather modifier, a monoalkyl or monoalkenyl phosphate salt (or mixtures thereof) having the structure (2):



20 where R^{O^3} is chosen from branched or unbranched alkyl or alkenyl groups having from 7 to 9 carbon atoms;

Y and Z are each chosen from H, alkali metal, ammonium and substituted ammonium counterions; and c is 0, or an integer of from 1 to 4.

25 Examples of the monoalkyl and monoalkenyl phosphate moiety include:

mono-n-heptyl phosphate
 mono-n-octyl phosphate
 mono-n-nonyl phosphate
 35 diethyleneglycol-mono-n-nonyl ether phosphate
 ethyleneglycol-mono-n-octyl ether phosphate
 mono-n-heptenyl phosphate
 mono-n-octenyl phosphate
 mono-n-nonenyl phosphate
 40 triethyleneglycol-mono-n-octenyl ether phosphate
 ethyleneglycol-mono-n-heptenyl ether phosphate
 mono-7-methyloctyl phosphate
 mono-5-methylhexyl phosphate
 mono-6,6-dimethylheptyl phosphate

45 Preferably, the R^2 group is the same as the R^1 group; thus when the dialkylphosphate salt is a dioctyl phosphate salt, then the monoalkyl phosphate is preferably a mono-octyl phosphate salt.

The amount of the monoalkyl or monoalkenyl phosphate salt which is present in the composition according to the invention is from 0.01 to 20%, preferably from 0.05 to 5% by weight of the composition.

50 Compositions containing less than 0.01% by weight of the monoalkyl or monoalkenyl phosphate salt are difficult to prepare, as the monosalt is difficult to separate entirely from the corresponding di-salt.

The weight ratio of the monoalkyl or monoalkenyl phosphate salt to the dialkyl or dialkenyl phosphate salt is from 1:1000 to 1:5, preferably from 1:1000 to 1:20.

Compositions in which this weight ratio is greater than 1:5 show poor lather characteristics.

55 The choice of an amount of the monoalkyl or monoalkenyl phosphate salt within the stated range will modify the lather, in terms of its volume and/or creaminess, that the composition is capable of producing. Evidence is given later in this specification to illustrate how the lather characteristics of the composition can be modified by varying the mono- and dialkyl or alkenyl:salt ratio.

Water

The composition according to the invention also comprises an amount of water to act as a vehicle for the dialkyl or dialkenyl phosphate salt and to enable it to be provided at a concentration suitable for convenient topical application to human skin.

The amount of water present in the composition of the invention is accordingly up to 99%, preferably from 5 to 99% by weight of the composition.

Co-surfactant

The composition according to the invention can also optionally comprise a co-surfactant, further to modify the surfactant properties attributable to the dialkyl or dialkenyl phosphate salt, and the monoalkyl or monoalkenyl phosphate salt.

Examples of co-surfactants include anionic surfactants other than the phosphate salts defined herein, as well as nonionic, amphoteric and zwitterionic surfactants.

Anionic Co-surfactants

Particularly preferred co-surfactants, when employed, are anionic surfactants, examples of which are set out hereinafter.

i. Fatty acid soap co-surfactant

The composition according to the invention can optionally comprise, as a co-surfactant one or more soaps which are water-soluble or water-dispersable alkali metal salts of an organic acid, especially a sodium or a potassium salt, or the corresponding ammonium or substituted ammonium salt. Examples of suitable organic acids are natural or synthetic alkanolic acids having from 10 to 22 carbon atoms, especially the fatty acids of triglyceride oils such as tallow and coconut oil.

For solid products, such as powders, bar or tablets, the preferred soap is a soap of tallow fatty acids, that is fatty acids derived from tallow class fats, for example beef tallow, mutton tallow, lard, palm oil and some vegetable butters. Minor amounts of up to about 30%, preferably 10 to 20%, by weight of sodium soaps of nut oil fatty acids derived from nut oils, for example coconut oil and palm kernel oil, may be admixed with the sodium tallow soaps, to improve their lathering and solubility characteristics if desired. Whereas tallow fatty acids are predominantly C₁₄ and C₁₈ fatty acids, the nut oil fatty acids are of shorter chain length and are predominantly C₁₀-C₁₄ fatty acids.

For liquid or gel products, the preferred soaps are predominantly C₁₀₋₁₄ fatty acids derived from nut oils, or alternatively, from synthetic alkanolic acids.

The soaps can be provided as a preformed ingredient for the composition, or they can be formed in situ during the manufacture of the composition by reaction of suitable fatty acids and an alkali.

The amount of fatty acid soap which can be present in the composition according to the invention is up to 90%, preferably from 2 to 80% by weight of the composition.

ii. Non-soap anionic co-surfactants

The composition according to the invention can also optionally comprise one or more non-soap anionic co-surfactants, examples of which include:

The alkali metal salts of organic sulphuric reaction products having an alkyl or acyl radical containing from 8-22 carbon atoms and a sulphonic acid or sulphuric acid ester group. Specific examples of these synthetic anionic surfactants are the sodium, ammonium, potassium or triethanolammonium alkyl sulphates, especially those obtained by sulphating the higher alcohols (C₈-C₁₈), sodium coconut oil fatty acid monoglyceride sulphates and sulphonates; sodium or potassium salts of sulphuric esters of the reaction product of one mole of a higher fatty alcohol (e.g. tallow or coconut oil alcohols) and 1-12 moles of ethyleneoxide; sodium or potassium salts of alkyl phenol ethylene oxide ether sulphate with 1-10 units of ethylene oxide per molecule and in which the alkyl group contains from 8 to 12 carbon atoms, sodium alkyl glyceryl ether sulphonates, the reaction product of fatty acids having from 10 to 22 carbon atoms esterified with isethionic acid and neutralised with sodium hydroxide; water soluble salts of condensation products of fatty acids with N-methyl taurine.

Especially preferred non-soap anionic co-surfactants include:

alkylaryl sulphonates, such as sodium alkyl benzene sulphonate (e.g. TEEPOL CM44, available from Shell).

alkyl sulphates, such as sodium lauryl sulphate (e.g. EMPICOL CX, available from Albright & Wilson), and


triethanolammonium lauryl sulphate (e.g. EMPICOL TL40/T, available from Albright & Wilson).

alkylether sulphonates, such as sodium lauryl ether sulphate (e.g. EMPICOL ESB70, available from Albright & Wilson).

alkyl sulphonates, such as sodium alkane (C13-18) sulphonate (e.g. HOSTAPUR SAS 30, available from Hoechst).

olefin sulphonates, such as sodium olefin sulphonate (C15-18) (e.g. HOSTAPUR OS, available from Hoechst).

Sarcosinates, having the structure (3):




where R³ is chosen from C₈₋₁₄ alkyl, and

M is a counterion chosen from alkali metals, ammonium, substituted ammonium, such as alkylolammonium.

An example of sarcosinates having the structure (3),

sodium lauryl sarcosinate (e.g. HAMPOSYL L-95, available from Grace).

Taurides, having the structure (4):




where R⁴ is chosen from C₈₋₁₈ alkyl

An example of taurides having the structure (4) is:

coconut methyl taurine (e.g. FENOPON TC 42, available from GAF).

Isethionates, having the structure (5):




where R⁵ is chosen from C₈₋₁₈ alkyl.

An example of isethionates having the structure (5) is:

sodium acyl isethionate (e.g. JORDAPON C1, available from Jordan).

Monoalkyl sulphosuccinates, having the structure (6):



where R^6 is chosen from C_{10-20} alkyl.

Examples of monoalkyl sulposuccinates having this structure (6) include:

sodium lauryl sulposuccinate (e.g. EMPICOL SLL, available from Albright & Wilson)

magnesium alkyl sulposuccinate (e.g. ELFANOL 616 Mg, available from AKZO),


sodium lauryl ethoxysulposuccinate (e.g. EMPICOL SDD, available from Albright & Wilson)

coconut monoethanolamide ethoxysulposuccinate, (e.g. EMPICOL SGG)

disodium lauryl polyglycoether sulposuccinate (e.g. SURTAGENE S30, available from CHEM-Y)

polyethyleneglycol sulposuccinate (e.g. REWOPOL SBFA 30, available from REWO).

Dialkyl sulposuccinates, having the structure (7):




where R^7 and R^8 are the same or different, and are chosen from C_{8-14} alkyl.

An example of dialkyl sulposuccinate having the structure (7) is:

sodium dioctyl sulposuccinate (e.g. EMCOL 4500 available from Witco).

Acyl lactylates, having the structure (8):



where R^9 is chosen from C_{8-18} alkyl,

and n is 1 or 2.

An example of acyl lactylates having the structure (8) is:

decanoyl lactylate (e.g. PATIONIC 122A, available from Patterson, C.J.).

Acylated α -amino acids, such as sodium lauroyl glutamate (e.g. ACYLGLUTAMATE LS-11, available from Ajinomoto Co. Inc.).

Ethyl carboxylates, such as alkyl $\text{C}_{12-14}\text{O}(\text{EO})_4\text{OCH}_2\text{CO}_2\text{Na}$ (e.g. AKYPO RLM 38, available from AKZO).

Nonionic co-surfactants


A composition according to the invention can also comprise nonionic co-surfactants which are compounds produced by the condensation of alkylene oxide groups (hydrophilic in nature) with an organic hydrophobic compound, which may be aliphatic or alkyl aromatic in nature. Examples of nonionic co-surfactants include:

- i. The polyethylene oxide condensates of alkyl phenols having from 8 to 12 carbon atoms, either straight or branched chain, with ethylene oxide, which is present in amounts of from 10 to 60 moles of ethylene

oxide per mole of alkylphenol.

ii. The condensation product of aliphatic alcohols having from 8 to 18 carbon atoms, straight or branched chain, with ethyleneoxide, for example, a coconut alcohol ethyleneoxide condensate having from 10 to 13 moles of ethylene oxide per mole of coconut alcohol, the coconut alcohol fraction having from 10 to 14 carbon atoms.

iii. Long chain tertiary amineoxides having the structure (9):



where R^{10} contains an alkyl, alkenyl or monohydroxyalkyl radical of from 8 to 18 carbon atoms, from 0 to 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety, and R^{11} and R^{12} contain from 1 to 3 carbon atoms and up to 1 hydroxy group, for example, methyl, ethyl, propyl, hydroxyethyl, or hydroxypropyl groups.

Especially preferred examples of nonionic cosurfactants include:

alkylethoxylates, such as the DOBANOL series, available from Shell;

esterethoxylates, such as the TAGAT series, available from Goldschmidt;

alkylalkanolamides, such as coconut monoethanolamide (e.g. EMPILAN CME, available from Albright & Wilson), and coconut diethanolamide, (e.g. EMPILAN CDE, available from Albright & Wilson).

sugar esters, such as sucrose laurate and methyl glucose laurate (available from Grillo-Werke A.G.)

esters of glycols, such as ethylene glycol mono stearate.

esters of glycerol, such as glyceryl mono stearate.

ethoxylated sorbitan esters, such as the TWEEN series (available from ICI).

amine oxides, such as alkyl dimethyl amine oxide (e.g. EMPIGEN OB, available from Albright & Wilson) and alkylethoxydimethyl amine oxide (e.g. EMPIGEN OY, available from Albright & Wilson).

Zwitterionic and Amphoteric co-surfactants

The composition according to the invention can also contain zwitterionic co-surfactants, which are derivatives of aliphatic quaternary ammonium, phosphonium and sulphonium compounds in which the aliphatic radicals can be straight or branched chain, and where one aliphatic substituent contains from 8 to 18 carbon atoms, and one contains an anionic water-solubilising group, such as carboxyl, sulphonate, sulphate, phosphate or phosphonate.

Examples of zwitterionic co-surfactant include:

4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate

5-N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxypentane-1-sulphate.

Particularly preferred zwitterionic co-surfactants are betaines, preferred examples of which are:

Alkyl betaines, having the structure (10):




where R^{16} is C_{10-18} alkyl.

An example of alkyl betaines having the structure (10) is:


lauryldimethyl betaine (e.g. EMPIGEN BB, available from Albright & Wilson).

Alkylamidopropyl betaines, having the structure (11):




where R¹⁷ is C₁₀₋₁₈ alkyl.

An example of alkylamidopropyl betaines having the structure (11) is:
cocamidopropyl betaine (e.g. TEGO BETAIN L7, available from Goldschmidt).
Alkylamphoglycinates, and having the structure (12):



where R¹⁸ is C₁₀₋₁₆ alkyl

R^{10} and R^{20} are the same or different and are chosen from H, CH_2COO^- and $(\text{CH}_2)_2\text{COO}^-$. An example of alkylamphoglycinates having the structure (12) is: cocoamphoglycinate (available from GAF), and alkoamphodipropionate. Sultaines, having the structure (13):



where R²¹ is chosen from C₁₂₋₁₈ alkyl or alkylamido.

An example of sultaines having the structure (13) is:
cocoamidopropylhydroxysultaine (e.g. CYCLOTERIC BET-CS, available from Alcolac).

Particularly preferred examples of amphoteric co-surfactants include:

Alkoamphoacetates, such as cocoamphoacetate (e.g. MIRANOL CM), and Alkoamphopropionates, such as cocoamphopropionate (e.g. MIRANOL CM-SF) both available from Miranol Inc.

The amount of co-surfactant when present in the compositions according to the invention is usually up to 50%, preferably from 1 to 40% by weight.

50 Optional thickening agent

The composition according to the invention can also comprise a polymer thickener in an amount sufficient to adjust the viscosity of the composition, so as to facilitate dispensing it conveniently onto the body surface.

Examples of polymer thickeners include:

anionic cellulose materials, such as sodium carboxy methyl cellulose;
anionic polymers such as carboxy vinyl polymers, for examples Carbomer 940 and 941;
nonionic cellulose materials, such a methyl cellulose and hydroxy propyl methyl cellulose;

cationic cellulose materials, such as Polymer JR 400;
 cationic gum materials, such as Jaguar C13 S;
 other gum materials such as gum acacia, gum tragacanth, locust bean gum, guar gum and carrageenan;
 proteins, such as albumin and protein hydrolysates;

5 and

clay materials, such as bentonite, hectorite, magnesium aluminium silicate, sodium magnesium silicate and a synthetic complex clay having the generic formula:

$[\text{Si}_6\text{Mg}_{6.1}\text{Li}_{6.8}\text{H}_{4.8}\text{O}_{24}]^{10.6-} \text{Na}^{+0.6}$, an example of which is Laponite, available from Laporte Industries.

10 The amount of thickening agent which can optionally be employed in the composition according to the invention is normally from 0.05 to 5%, preferably 0.1 to 1% by weight of the composition.

Preservative

15 The composition according to the invention can also optionally comprise a preservative to prevent microbial spoilage, especially biodegradation of the alkyl phosphate salt. It is accordingly apparent that the composition containing the alkyl phosphate salt may be prone to attack by bacteria, moulds and fungi and other microbial influences. There is therefore a risk that the shelf-life of the composition might be unacceptably short due to the biodegradation or spoilage, unless there is included in the composition a bactericide, fungicide or other microbicide in an amount sufficient to inhibit or prevent the said biodegradation or spoilage, or unless other steps
 20 are taken to preserve the composition.

Examples of preservatives include:

- 25 (i) Chemical preservatives, such as ethanol, benzoic acid, sodium benzoate, sorbic acid, potassium sorbate, sodium propionate and the methyl, ethyl, propyl and butyl esters of p-hydroxybenzoic acid 2-bromo-2-nitropropane-1,3-diol, phenoxyethanol, dibromodicyanobutane, formalin and Triclosan. The amount of chemical preservative optionally to be incorporated in the composition according to the invention will generally be from 0.05 to 5%, preferably from 0.1 to 2% by weight, the amount chosen being sufficient to arrest microbial proliferation.
- 30 (ii) Water activity depressants, such as glycerol, propylene glycol, sorbitol, sugars and salts, for examples alkali metal halides, sulphates and carboxylates. When employing a water activity depressant, sufficient should be incorporated in the composition according to the invention to reduce the water activity (a_w) from 1 to <0.9, preferably to <0.85 and most preferably < 0.8, the lowest of these values being that at which yeasts, moulds and fungi will not proliferate.

35

Further optional ingredients

40 The composition according to the invention can also contain other optional adjuncts, that is ingredients other than the main ingredients already defined which are conventionally employed in compositions for topical application to human skin. These adjuncts, when present, will normally form the balance of the composition.

Examples of optional adjuncts include vehicles, the selection of which will depend on the required product form of the composition. Typically, the vehicle when present, will be chosen from diluents, dispersants or carriers for the dialkyl or dialkenyl phosphate salt so as to ensure an even distribution of it when applied to the skin.

45

Compositions according to this invention can include water as a vehicle, usually with at least one other cosmetically-acceptable vehicle.

Vehicles other than water that can be used in compositions according to the invention can include liquids or solids as emollients, solvents, humectants, thickeners and powders. Examples of each of these types of vehicles, which can be used singly or as mixtures of one or more vehicles, are as follows:

50

Emollients, such as stearyl alcohol, glyceryl monolaurate, glyceryl monoricinoleate, glyceryl monostearate, propane-1,2-diol, butane-1,3-diol, docosan-1,2-diol, mink oil, cetyl alcohol, isopropyl isostearate, stearic acid, isobutyl palmitate, isocetyl stearate, oleyl alcohol, isopropyl laurate, hexyl laurate, decyl oleate, octadecan-2-ol, isocetyl alcohol, eicosanyl alcohol, behenyl alcohol, cetyl palmitate, silicone oils such as dimethylpolysiloxane, di-n-butyl sebacate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, butyl stearate, polythylene glycol, triethylene glycol, lanolin, cocoa butter, corn oil, cotton seed oil, tallow, lard, olive oil, palm kernel oil, rapeseed oil, safflower seed oil, soybean oil, sunflower seed oil, olive oil, sesame seed oil, coconut oil, arachis oil, castor oil, acetylated lanolin alcohols, petroleum, mineral oil, butyl myristate, isostearic acid, palmitic acid, isopropyl linoleate, lauryl lactate, myristyl lactate, decyl oleate, myristyl myristate;

55

Propellants, such as trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethane, monochlorodifluoromethane, trichlorotrifluoroethane, propane, butane, isobutane, dimethyl ether, carbon dioxide, nitrous oxide;

5 Solvents, such as ethyl alcohol, methylene chloride, isopropanol, acetone, castor oil, ethylene glycol monoethyl ether, diethylene glycol monobutyl ether, diethylene glycol monoethyl ether, dimethyl sulphoxide, dimethyl formamide, tetrahydrofuran;

Humectants, such as glycerin, sorbitol, sodium 2-pyrrolidone-5-carboxylate, soluble collagen, dibutyl phthalate, gelatin;

10 Powders, such as chalk, talc, fullers earth, kaolin, starch, gums, colloidal silicon dioxide, sodium polyacrylate, tetra alkyl and/or trialkyl aryl ammonium smectites, chemically modified magnesium aluminium silicate, organically modified montmorillonite clay, hydrated aluminium silicate, fumed silica, carboxyvinyl polymer, sodium carboxymethyl cellulose, ethylene glycol monostearate.

The cosmetically acceptable vehicle, when present, will usually form from 0.01 to 99.9%, preferably from 50 to 98% by weight of the composition, and can, in the absence of other cosmetic adjuncts, form the balance of the composition.

PRODUCT FORM OF THE COMPOSITION

20 The composition according to the invention can take the form of a liquid or gel, intended to be dispensed from a capped container such as a bottle, roll-on applicator or tube, or a pump-operated or propellant-driven aerosol dispenser, as a skin cleanser, shower product, bath additive or shampoo. The composition can also take the form of a powder or a solid such as a stick, preferably housed in a suitable capped holder with a wind-up or push-up action similar to a lip stick, or a bar or tablet, with or without fatty acid soaps, intended to be used for washing instead of a conventional soap bar.

25 The invention also provides a closed container containing a detergent composition as herein defined.

Process for Preparing the Composition

30 The invention also provides a process for preparing the composition of the type defined herein, which process comprises the steps of:

- (i) preparing a mixture comprising one or more dialkyl and/or dialkenyl phosphate salts, as defined herein, and one or more monoalkyl and/or monoalkenyl phosphate salts as defined herein, and
- (ii) subsequently packaging the mixture into containers.

Use of the composition

The composition according to the invention is intended primarily as a personal washing product for cleansing the face and other sensitive parts of the body surface, including the mucosae. It can also be used for washing the hair as well as the skin.

40 In use, a small quantity, for example from 1 to 5 ml, of the composition is either rubbed between the hands, together with water to form a foam, which is then used for washing, or applied via a flannel or sponge to the area to be cleansed, or the foam is generated directly on that area. The foam is subsequently rinsed away with clean water.

EVIDENCE TO SHOW ENHANCED LATHER PROPERTIES

Experiments were performed to demonstrate how lather volume obtainable with dioctyl phosphate salt increases with decreasing levels of monoethyl phosphate salt in compositions according to the invention.

50 Foam was generated in a standard manner using a Foam Machine whose construction and use will now be described.

The Foam Machine

55 Foam is generated within a glass graduated cylinder by the action of a perforated piston upon the test composition and a suitably treated substrate. The repetitive compression and expansion of the substrate by the piston results in the aeration of the composition and the generation of foam which accumulates above the piston plate. The volume of the foam can then be read using the graduations of the cylinder when the piston is at its lowest point. The apparatus is enclosed and is supplied with air at a constant temperature of 38°C.

The operation of the machine in detail is as follows. Crimped nylon fibre is soaked in a large quantity of a solution of 2.5% by weight artificial sebum, consisting of squalene, cholesterol, long chain fatty acids, long chain triglycerides and decyl oleate, in dichloromethane for 1 hour. The solvent is poured away, and the fibres are dried by evaporation of residual solvent. The sebum-treated nylon fibre is intended to simulate greasy human skin or hair to which the composition of the invention would normally be applied when washing the skin or shampooing the hair.

Into a graduated glass cylinder of internal diameter 3.8cm are placed two discs of polyurethane foam (diameter 3.8cm, height 1.0cm). 2g of sebum-treated fibre is placed on top of the foam and the diluted test composition (2 ml) is poured over. The diluted composition is prepared from 7.5g of test composition and 92.5g of distilled water.

The prepared cylinder is placed within the foam machine, and with the cross-head at its lowest point the piston plate is adjusted to a level of 70 ml (as determined from the graduations on the cylinder).

The machine is operated with a periodicity of 12 rpm. After each minute of operation, the machine is stopped with the cross-head at the highest point and the volume of foam above the piston plate is recorded.

The machine is operated for a total of 10 minutes, the reported value is the mean of the foam volume values recorded between 3 and 10 minutes inclusive.

Foam Volume Results

All foam values for the dialkyl phosphate were recorded for the triethanolamine salt at pH 8. The percentage surfactant in the composition is based on the weight of the dialkyl phosphate in the acid form, not as the salt.

The following table shows the Foam Volumes obtainable from an aqueous solution containing dioctyl phosphate (DOP) and mono-octyl phosphate (MOP) at a range of values.

DOP (% w/w)	MOP (% w/w)	Foam Volume (ml)
20	0	54.3
19	1	38.9
18	2	26.3
17	3	29.4

From these results, it can be seen that the foam volume is enhanced with decreasing levels of mono-octyl phosphate in relation to the amounts of dioctyl phosphate present. Preferably the weight ratio of mono-octyl phosphate to dioctyl phosphate should not exceed 3:17.

EXAMPLES

The invention is further illustrated by reference to the following examples.

Example 1

This Example illustrates a body cleansing liquid product suitable for use in the shower.

The product contains the following ingredients:

	<u>Ingredients</u>	<u>% w/w</u>
5	triethanolamine di-n-octyl phosphate	36
	triethanolamine mono-n-octyl phosphate	0.1
	preservative	2
10	water	to 100

This product can be used for cleansing the whole body surface, including the hair, for example under the shower, a convenient amount of say 5ml being placed in the palm of the hand prior to distributing over the body surface with added water to create a lather with superior volume and creaminess characteristics.

15 Examples 2 to 8 illustrate body shampoos for use in the shower or when bathing.

	<u>Example 2</u>	<u>% w/w</u>
20	triethanolammonium di-n-octyl phosphate	36
	triethanolammonium mono-n-octyl phosphate	0.2
25	triethanolammonium laurate	1
	triethanolammonium myristate	2
	alkyldialkylamine oxide (EMPIGEN OB:	
	30% active	8.3
30	cationised cellulose	0.1
	propylene glycol	10
	water	to 100
35		

	<u>Example 3</u>	<u>% w/w</u>
40	triethanolammonium di-n-octyl phosphate	24
	triethanolammonium mono-n-octyl phosphate	0.2
45	lauryl dimethylbetaine	3
	ethylene glycol monostearate	1.5
	propylene glycol	2.5
50	preservative, perfume, dyes	q.v.
	water	to 100%
55		

Example 4

5		<u>% w/w</u>
	sodium di-(diethyleneglycol-mono-n-nonyl	
	ether) phosphate	30
10	sodium mono-diethyleneglycol-mono-n-nonyl	
	ether) phosphate	0.4
	triethanolammonium laurate	1
15	triethanolammonium myristate	2
	alkyldialkylamine oxide (EMIGEN OB:	
	30% active)	8.3
	cationised cellulose	0.1
20	propylene glycol	10
	water	to 100

Example 5

		<u>% w/w</u>
30	triethanolammonium di-(ethyleneglycol	
	-mono-n-octyl ether) phosphate	20
35	triethanolammonium mono-(ethyleneglycol	
	-mono-n-octyl ether) phosphate	2
	lauryl dimethylbetaine	3
	ethylene glycol monostearate	1.5
40	propylene glycol	2.5
	preservative, perfume, dyes	q.v.
	water	to 100%

Example 6

5		<u>% w/w</u>
	triethanolammoniumdi-(ethyleneglycol	
	-mono-n-heptyl ether) phosphate	30
10	triethanolammonium mono-(ethyleneglycol	
	-n-heptyl ether) phosphate	0.2
	triethanolammonium laurate	1
15	triethanolammonium myristate	2
	alkyldialkylamine oxide (EMIGEN OB:	
	30% active)	8.3
	cationised cellulose	0.1
20	propylene glycol	10
	water	to 100

Example 7

25		<u>% w/w</u>
30	sodium di-(7-methyloctyl) phosphate	25
	sodium mono-(7-methyloctyl) phosphate	0.3
35	lauryl dimethylbetaine	3
	ethylene glycol monostearate	1.5
	propylene glycol	2.5
	preservative, perfume, dyes	q.v.
40	water	to 100%

g w/wExample 8

5	sodium di-(di-ethyleneglycol-mono-n-nonyl ether) phosphate	25
10	sodium mono-(ethyleneglycol-mono-n-nonyl ether) phosphate	0.5
	cocoamidopropyl betaine	5
	preservatives, dyes	q.v.
15	water	to 100

Examples 9 to 16 illustrate hair shampoos.

g w/wExample 9

25	triethanolammonium di-n-heptyl phosphate	20
	triethanolammonium mono-n-heptyl phosphate	0.5
30	cocamidopropyl betaine	1.5
	preservatives, dyes, perfumes	q.v.
	water	to 100%

Example 10

40	triethanolammonium di-n-nonyl phosphate	10
	disodium lauryl sulphosuccinate	4
	cocamidopropyl betaine	4
	triethanolammonium mono-n-nonyl phosphate	0.8
45	preservatives, dyes, perfumes	q.v.
	water	to 100%

% w/wExample 11

5

triethanolammonium di-n-heptenyl phosphate	20
triethanolammonium mono-n-heptenyl phosphate	0.5
10 cocamidopropyl betaine	1.5
preservatives, dyes, perfumes	q.v.
water	to 100%

15

Example 12

20

% w/w

triethanolammonium di-n-nonenyl phosphate	10
triethanolammonium mono-n-nonenyl phosphate	0.8
25 disodium lauryl sulphosuccinate	4
cocamidopropyl betaine	4
preservatives, dyes, perfumes	q.v.
30 water	to 100%

35

% w/wExample 13

40

potassium di-(5-methylhexyl) phosphate	25
potassium mono-(5-methylhexyl) phosphate	0.5
cocamidopropyl betaine	1.5
45 preservatives, dyes, perfumes	q.v.
water	to 100%

50

55

Example 14

5	triethanolammonium di-(6,6-dimethylheptyl)	
	phosphate	10
	disodium lauryl sulphosuccinate	4
10	cocamidopropyl betaine	4
	triethanolammonium mono-	
	(6,6-dimethylheptyl) phosphate	1
15	preservatives, dyes, perfumes	q.v.
	water	to 100%

Example 15

		<u>% w/w</u>
25	triethanolammonium di-octenyl phosphate	15
	triethanolammonium mono-octenyl phosphate	0.3
	lauryldimethyl betaine	6
	preservatives, perfumes, dyes	q.v.
30	water	to 100

Example 16

		<u>% w/w</u>
40	triethanolammonium di-(7-methyloctyl) phosphate	25
	triethanolammonium mono-(7-methyloctyl) phosphate	0.4
	preservatives, perfumes, dyes	q.v.
	water	to 100

45 Examples 17 to 21 illustrate facial foaming cleansers.

50

55

% w/wExample 17

5		
	sodium di-n-octylphosphate	30
	sodium mono-n-octyl phosphate	0.6
10	lauryl dimethyl betaine	4
	sorbitol	5
	glycerol	2
	quaternised cellulosic polymer (Polymer	
15	JR400)	0.3
	preservative, dyes, perfumes	q.v.
	water	to 100

20

% w/wExample 18

25		
	sodium di-(triethyleneglycol-mono-n-octenyl	
	ether) phosphate	25
	sodium mono-(triethyleneglycol-n-octenyl	
30	ether) phosphate	0.5
	lauryl dimethyl betaine	4
	sorbitol	5
35	glycerol	2
	quaternised cellulosic polymer (Polymer	
	JR400)	0.3
	preservative, dyes, perfumes	q.v.
40	water	to 100

45

50

55

	<u>% w/w</u>
5 <u>Example 19</u>	
sodium di-n-octyl phosphate	30
sodium mono-n-octyl phosphate	0.6
10 lauryl dimethyl betaine	4
sorbitol	5
glycerol	2
15 quaternised cellulosic polymer (Polymer JR400)	0.3
preservative, dyes, perfumes	q.v.
20 water	to 100

	<u>% w/w</u>
25 <u>Example 20</u>	
sodium di-n-octylphosphate	15
sodium di-n-nonylphosphate	10
30 sodium mono-n-octyl phosphate	0.1
sodium mono-n-nonyl phosphate	0.1
sorbitol	3
35 preservatives, perfume, dyes	q.v.
water	to 100%

40 <u>Example 21</u>	
sodium di-(diethyleneglycol-mono-n-octyl 45 ether) phosphate	30
sodium mono-(diethyleneglycol-mono-n-octyl ether) phosphate	0.8
50 sorbitol	6
propylene glycol	2
preservatives, perfumes, dyes	q.v.
55 water	to 100

Examples 22, 23 & 24

The following three examples illustrate soap-containing products with di-n-octyl phosphate salt and mono-n-octyl phosphate in accordance with the invention.


In each case, the products were made in accordance with standard soap manufacture. The product of Example 22 yielded a flaked product which could not readily be pressed into a bar, as the flakes were not sufficiently cohesive. With the products of Examples 23 and 24 bars were formed following the usual plodding, extrusion and stamping that is conventional in soap bar manufacture.

These soap products had the following formulations:


Ingredient	Examples		
	22	23	24
	(% by weight)		
Hardened Tallow Soap	64	66.6	-
80/20 tallow-coco soap	-	9	76
sodium di-n-octyl phosphate	15.8	14.3	18.8
sodium mono-n-octyl phosphate	0.2	0.1	0.2
water	20	10	5

Claims

1. A composition suitable for topical application to the skin or hair, which comprises:
- from 1 to 99.9% by weight of a dialkyl or dialkenyl phosphate salt (or mixtures thereof) having the structure (1):



- from 0.01 to 20% by weight of a monoalkyl or monoalkenyl phosphate salt (or mixtures thereof) having the structure (2):



where R^{01} , R^{02} and R^{03} are each chosen from branched or unbranched, alkyl or alkenyl groups having an average of from 7 to 9 carbon atoms;

X, Y and Z are each chosen from H, alkali metal, ammonium and substituted ammonium counterions; and

5 a, b and c are each chosen from 0 or a value of from 1 to 4;

the weight ratio of the monoalkyl or monoalkenyl phosphate salt to the dialkyl or dialkenyl phosphate salt being from 1:1000 to 1:5.

2. A composition according to claim 1, in which the dialkyl phosphate moiety of the alkyl phosphate salt is chosen from:

10 di-n-heptyl phosphate
di-n-octyl phosphate
di-n-nonyl phosphate
di-(7-methyloctyl) phosphate
15 di-(5-methylhexyl) phosphate
di-(6,8-dimethylheptyl) phosphate
di-(ethyleneglycol-mono-n-octyl ether) phosphate
di-(ethyleneglycol-mono-n-heptyl ether) phosphate
di-(diethyleneglycol-mono-n-octyl ether) phosphate
20 n-hexyl-n-nonyl phosphate, and
n-hexyl-n-octyl phosphate.

3. A composition according to claim 1, in which the dialkenyl phosphate moiety of the alkenyl phosphate salt is chosen from:

25 di-n-heptenyl phosphate
di-n-octenyl phosphate
di-n-nonenyl phosphate, and
di-(triethyleneglycol-mono-n-octenyl ether) phosphate

4. A composition according to claim 1 or 2, in which the dialkyl phosphate salt is a dioctyl phosphate salt.

5. A composition according to any preceding claim, in which the monoalkyl phosphate moiety of the monoalkyl phosphate salt is chosen from:

35 mono-n-heptyl phosphate
mono-n-octyl phosphate
mono-n-nonyl phosphate
diethyleneglycol-mono-n-nonyl ether phosphate
ethyleneglycol-mono-n-octyl ether phosphate
mono-7-methyloctyl phosphate, and
40 mono-6,8-dimethylheptyl phosphate

6. A composition according to any of claims 1 to 6, in which the monoalkenyl phosphate moiety of the monoalkenyl phosphate salt is chosen from:

45 mono-n-heptenyl phosphate
mono-n-octenyl phosphate
mono-n-nonenyl phosphate, and
triethyleneglycol-mono-n-octenyl ether phosphate


7. A composition according to claims 5 or 6, in which the monoalkenyl phosphate salt is a mono-n-octenyl phosphate salt.

8. A composition according to any preceding claim, in which the counterion is chosen from sodium, potassium or triethanolammonium.


9. A composition according to any of claims 1 to 4 in which the dialkyl or dialkenyl phosphate salt forms from 2 to 50% by weight of the composition.

10. A composition according to any of claims 1 and 5 to 7 in which the monoalkyl or monoalkenyl phosphate salt forms from 0.05 to 5% by weight of the composition.

11. A composition according to any preceding claim, in which the alkyl group of the monoalkyl or monoalkenyl phosphate salt has the same number of carbon atoms as that of the dialkyl or dialkenyl group of the phosphate salt.
12. A composition according to any preceding claim, in which the weight ratio of the monoalkyl or monoalkenyl phosphate salt to the dialkyl or dialkenyl phosphate salt is from 1:1000 to 1:20.
13. A composition according to any preceding claim, which further comprises an anionic co-surfactant.
14. A composition according to any preceding claim, which further comprises a nonionic co-surfactant.
15. A composition according to any preceding claim, which further comprises a zwitterionic co-surfactant.
16. A composition according to any preceding claim, which is a liquid or gel product.
17. A composition according to claim 16, which is a washing product suitable for topical application to the face.
18. A composition according to any of claims 1 to 16, which is a shampoo.
19. A composition according to any of claims 1 to 15, which is a powder.
20. A composition according to any of claims 1 to 15, which is a bar or tablet suitable for washing the skin.
21. A composition according to claim 20, which is a soap bar or soap tablet.
22. A process for preparing a composition according to any preceding claim, which process comprises the steps of:
- preparing a mixture comprising:
 - one or more dialkyl or dialkenyl phosphate salts having the structure (1):



- one or more monoalkyl or monoalkenyl phosphate salts having the structure (2):



where R^{01} , R^{02} and R^{03} are each chosen from branched or unbranched, alkyl or alkenyl groups having an average of from 7 to 9 carbon atoms; and

X, Y and Z are each chosen from H, alkali metal, ammonium, substituted ammonium counterions; and

a, b and c are each chosen from 0, or a value of from 1 to 4;


the weight ratio of the monoalkyl or monoalkenyl phosphate salt to the dialkyl or dialkenyl phosphate salt being from 1:1000 to 1:5, and

ii. subsequently packaging the composition so formed into containers.


23. The use of a composition according to any of claims 1 to 21 for washing human skin or hair.

Patentanaprüche

1. Zusammensetzung, geeignet für topische Anwendung auf die Haut oder das Haar, welche umfaßt
a. von 1 bis 99,9 Gewichtsprozent eines Dialkyl- oder Dialkenylphosphatsalzes (oder Mischungen dar-
aus) mit der Struktur (1):



- b. von 0,01 bis 20 Gewichtsprozent eines Monoalkyl- oder Monoalkenylphosphatsalzes (oder Mischungen daraus) der Struktur (2):



worin jeder der Reste R^{01} , R^{02} und R^{03} aus verzweigten oder unverzweigten Alkyl- oder Alkenylgruppen mit einer durchschnittlichen Zahl von 7 bis 9 Kohlenstoffatomen ausgewählt ist,

jeder der Reste X, Y und Z aus H-, Alkalimetall-, Ammonium- und substituierten Ammonium-Genen ausgewählt ist, und

a, b und c jeweils den Wert 0 oder einen Wert von 1 bis 4 besitzen,

wobei das Gewichtsverhältnis des Monoalkyl- oder Monoalkenylphosphatsalzes zu dem Dialkyl- oder Dialkenylphosphatsalz im Bereich von 1:1000 bis 1:5 liegt.

2. Zusammensetzung gemäß Anspruch 1, in welcher der Dialkylphosphat-Teil des Alkylphosphatsalzes ausgewählt ist aus:

Di-n-heptylphosphat,
Di-n-octylphosphat,
Di-n-nonylphosphat,
Di-(7-methyloctyl)-phosphat,
Di-(5-methylhexyl)-phosphat,
Di-(6,6-dimethylheptyl)-phosphat,
Di-(ethylenglykol-mono-n-octylether)-phosphat,
Di-(ethylenglykol-mono-n-heptylether)-phosphat,
Di-(diethylenglykol-mono-n-octylether)-phosphat,
n-Hexyl-n-nonylphosphat und
n-Hexyl-n-octylphosphat.

3. Zusammensetzung gemäß Anspruch 1, in welcher der Dialkenylphosphat-Teil des Alkenylphosphatsalzes ausgewählt ist aus:

Di-n-heptenylphosphat,
Di-n-octenylphosphat,
Di-n-nonenylphosphat und
Di-(triethylenglykol-mono-n-octenylether)-phosphat.


4. Zusammensetzung gemäß einem der Ansprüche 1 oder 2, in welcher das Dialkylphosphatsalz ein Dioctylphosphatsalz ist.

5. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, in welcher der Monoalkylphosphat-Teil des Monoalkylphosphatsalzes ausgewählt ist aus:


Mono-n-heptylphosphat,
Mono-n-octylphosphat,
Mono-n-nonylphosphat,

Diethylenglykol-mono-n-nonylether-phosphat,
Ethylenglykol-mono-n-octylether-phosphat,
Mono-7-methyloctylphosphat und
Mono-6,6-dimethylheptylphosphat.

- 5 6. Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 6, in welcher der Monoalkenylphosphat-Teil des Monoalkenylphosphatsalzes ausgewählt ist aus:
Mono-n-heptenylphosphat,
Mono-n-octenylphosphat,
10 Mono-n-nonenylphosphat und
Triethylenglykol-mono-n-octenylether-phosphat.
7. Zusammensetzung gemäß irgendeinem der Ansprüche 5 oder 6, in welcher das Monoalkenylphosphatsalz ein Mono-n-octenylphosphatsalz ist.
- 15 8. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, in welcher das Gegenion aus Natrium, Kalium oder Triethanolammonium ausgewählt ist.
9. Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 4, in welcher das Dialkyl- oder Dialkenylphosphatsalz von 2 bis 50 Gewichtsprozent der Zusammensetzung ausmacht.
- 20 10. Zusammensetzung gemäß irgendeinem der Ansprüche 1 und 5 bis 7, in welcher das Monoalkyl- oder Monoalkenylphosphatsalz von 0,05 bis 5 Gewichtsprozent der Zusammensetzung ausmacht.
11. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, in welcher die Alkylgruppe des Monoalkyl- oder Monoalkenylphosphatsalzes die gleiche Anzahl von Kohlenstoffatomen aufweist, wie die der Dialkyl- oder Dialkenylgruppe des Phosphatsalzes.
- 25 12. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, in welcher das Gewichtsverhältnis des Monoalkyl- oder Monoalkenylphosphatsalzes zu dem Dialkyl- oder Dialkenylphosphatsalz im Bereich von 1:1000 bis 1:20 liegt.
- 30 13. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, welche ferner noch ein anionisches Hilfssurfactant enthält.
14. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, welche ferner noch ein nichtionisches Hilfssurfactant enthält.
- 35 15. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, welche ferner noch ein zwitterionisches Hilfssurfactant enthält.
- 40 16. Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, welche ein Flüssig- oder Gelprodukt ist.
17. Zusammensetzung gemäß Anspruch 16, die ein für topische Anwendung auf das Gesicht geeignetes Waschprodukt ist.
- 45 18. Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 16, welche ein Shampoo ist.
19. Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 15, welche ein Pulver ist.
20. Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 15, welche ein Riegel oder eine Tablette ist, die für das Waschen der Haut geeignet sind.
- 50 21. Zusammensetzung gemäß Anspruch 20, welche ein Seifenriegel oder eine Seifentablette ist.
22. Verfahren zur Herstellung einer Zusammensetzung gemäß irgendeinem der vorhergehenden Ansprüche, welches die Stufen umfaßt:
55 I. Herstellung einer Mischung enthaltend:
a. ein oder mehrere Dialkyl- oder Dialkenylphosphatsalze der Struktur (1):



b. ein oder mehrere Monoalkyl- oder Monoalkenylphosphatsalze der Struktur (2):



worin jeder der Reste R^{01} , R^{02} und R^{03} aus verzweigten oder unverzweigten Alkyl- oder Alkenylgruppen mit einer durchschnittlichen Zahl von 7 bis 9 Kohlenstoffatomen ausgewählt ist, und jeder der Reste X, Y und Z aus H-, Alkalimetall-, Ammonium-, substituierten Ammonium-Gegenionen ausgewählt ist, und

a, b und c jeweils den Wert 0 oder einen Wert von 1 bis 4 besitzen,


wobei das Gewichtsverhältnis des Monoalkyl- oder Monoalkenylphosphatsalzes zu dem Dialkyl- oder Dialkenylphosphatsalz im Bereich von 1:1000 bis 1:5 liegt und

ii. anschließendes Verpacken der so gebildeten Zusammensetzung in Behälter.


23. Die Verwendung einer Zusammensetzung gemäß irgendeinem der Ansprüche 1 bis 21 zum Waschen der menschlichen Haut oder des Haars.

Revendications

1. Une composition appropriée à une application topique sur la peau ou la chevelure, qui comprend :
a. de 1 à 99,9% en poids d'un sel de dialcényle ou de dialcyle phosphate (ou des mélanges de ceux-ci) ayant la structure (1) :



b. de 0,01 à 20% en poids d'un sel de monoalkyle ou monoalkényle phosphate (ou des mélanges de ceux-ci) ayant la structure (2) :



où R^{01} , R^{02} et R^{03} sont chacun choisis parmi des groupements alkyle ou alkényle, ramifiés ou non ramifiés ayant une moyenne de 7 à 9 atomes de carbone;


X, Y et Z sont chacun choisis parmi les contre-ions H, métal alcalin, ammonium et ammonium substitués; et a, b et c sont chacun choisis entre 0 ou une valeur allant de 1 à 4;

le rapport pondéral du sel de monoalkyle ou monoalkényle phosphate au sel de dialcyle ou


dialkényle phosphate allant de 1:1000 à 1:5.

2. Une composition selon la revendication 1, dans laquelle la partie dialkyle phosphate du sel d'alkyle phosphate est choisi parmi les :
 - 5 di-n-heptyle phosphate
 - di-n-octyle phosphate
 - di-n-nonyle phosphate
 - di-(7-méthyl-octyle) phosphate
 - di-(5-méthyl-hexyle) phosphate
 - 10 di-(6,6-diméthyl-heptyle) phosphate
 - di-(éther éthylèneglycol-mono-n-octylique) phosphate
 - di-(éther éthylèneglycol-mono-n-heptylique) phosphate
 - di-(éther diéthylèneglycol-mono-n-octylique) phosphate
 - n-hexyle-n-nonyle phosphate, et
 - 15 n-hexyle-n-octyle phosphate.
3. Une composition selon la revendication 1, dans laquelle la partie dialkylényle phosphate du sel d'alkényle phosphate est choisi parmi les :
 - di-n-heptényle phosphate
 - 20 di-n-octényle phosphate
 - di-n-nonényle phosphate, et
 - di-(éther triéthylèneglycol-mono-n-octénylique) phosphate
4. Une composition selon la revendication 1 ou 2, dans laquelle le sel de dialkyle phosphate est un sel de dioctyle phosphate.
5. Une composition selon l'une quelconque des revendications précédentes, dans laquelle la partie monoalkyle phosphate du sel de monoalkyle phosphate est choisi parmi les :
 - mono-n-heptyle phosphate
 - mono-n-octyle phosphate
 - 30 mono-n-nonyle phosphate
 - éther diéthylèneglycol-mono-n-nonylique phosphate
 - éther éthylèneglycol-mono-n-octylique phosphate
 - mono-7-méthyl-octyle phosphate, et
 - mono-6,6-diméthyl-heptyle phosphate
6. Une composition selon l'une quelconque des revendications 1 à 6, dans laquelle la partie monoalkylényle phosphate du sel de monoalkényle phosphate est choisi parmi les :
 - mono-n-heptényle phosphate
 - mono-n-octényle phosphate
 - 40 mono-n-nonényle phosphate, et
 - éther triéthylèneglycol-mono-n-octénylique phosphate
7. Une composition selon les revendications 5 ou 6, dans laquelle le sel de monoalkényle phosphate est un sel de mono-n-octényle phosphate.
8. Une composition selon l'une quelconque des revendications précédentes, dans laquelle le contre-ion est choisi parmi le sodium, le potassium ou le triéthanolammonium.
9. Une composition selon l'une des revendications 1 à 4 dans laquelle le sel de dialkyle ou dialkényle phosphate constitue de 2 à 50% en poids de la composition.
10. Une composition selon l'une quelconque des revendications 1 et 5 à 7 dans laquelle le sel de monoalkyle ou monoalkényle phosphate forme de 0,05 à 5% en poids de la composition.
11. Une composition selon l'une quelconque des revendications précédentes, dans laquelle le groupement alkyle du sel de monoalkyle ou monoalkényle phosphate a le même nombre d'atomes de carbone que celui du groupement dialkyle ou dialkényle du sel de phosphate.

12. Une composition selon l'une quelconque des revendications précédentes, dans laquelle le rapport en masse du sel de monoalkyle ou monoalkényle phosphate sur le sel de dialkyle ou dialkényle phosphate va de 1:1000 à 1:20.
13. Une composition selon l'une quelconque des revendications précédentes, qui comprend en plus un co-surfactant anionique.
14. Une composition selon l'une quelconque des revendications précédentes, qui comprend en plus un co-surfactant non-ionique.
15. Une composition selon l'une quelconque des revendications précédentes, qui comprend en plus un co-surfactant zwitterionique.
16. Une composition selon l'une quelconque des revendications précédentes, qui est un produit liquide ou en gel.
17. Une composition selon la revendication 16, qui est un produit de toilette convenant à une application topique sur le visage.
18. Une composition selon l'une quelconque des revendications 1 à 16, qui est un shampoing.
19. Une composition selon l'une quelconque des revendications 1 à 15, qui est une poudre.
20. Une composition selon l'une quelconque des revendications 1 à 15, qui est un pain ou une tablette appropriée pour la toilette de la peau.
21. Une composition selon la revendication 20, qui est un savon ou une savonnette.
22. Un procédé pour préparer une composition selon l'une quelconque des revendications précédentes, lequel procédé comprend les étapes de :
 - i. préparer un mélange comprenant
 - a. un ou plusieurs sels de dialkényle ou de dialkyle phosphate ayant la structure (1) :



- b. un ou plusieurs sels de monoalkyle ou monoalkényle phosphate ayant la structure (2) :



où R^{O1}, R^{O2} et R^{O3} sont chacun choisis parmi des groupements alkyle ou alkényle, ramifiés ou non ramifiés ayant une moyenne de 7 à 9 atomes de carbone;

X, Y et Z sont chacun choisis parmi les contre-ions H, métal alcalin, ammonium et ammonium substitués; et

a, b et c sont chacun choisis entre 0 ou une valeur allant de 1 à 4;

le rapport pondéral du sel de monoalkyle ou monoalkényle phosphate au sel de dialkyle ou dialkényle phosphate allant de 1:1000 à 1:5, et

ii. conditionner ensuite la composition ainsi formée dans des récipients.

23. L'utilisation d'une composition selon l'une quelconque des revendications 1 à 21 pour laver la peau ou la chevelure humaine.

5

10

15

20

25

30

35

40

45

50

55